



***Salmonella* spp., *Vibrio* spp., *Clostridium perfringens*, and *Plesiomonas shigelloides* in marine and freshwater invertebrates from coastal California ecosystems.**

Journal:	Microbial Ecology
Manuscript ID:	MECO-2005-0230.R1
Manuscript Type:	original manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	Miller, Woutrina Miller, Melissa Gardner, Ian Atwill, Rob Byrne, Barbara Jang, Spencer Harris, Michael Ames, Jack Jessup, Dave Paradies, Dave Worcester, Karen Melli, Ann Conrad, Patricia
Key Words:	fecal bacteria, invertebrate, salmonella, vibrio, clostridium

powered by ScholarOne
Manuscript Central™

Salmonella spp., *Vibrio* spp., *Clostridium perfringens*, and *Plesiomonas shigelloides* in marine and freshwater invertebrates from coastal California ecosystems.

W.A. Miller¹, M.A. Miller², I.A. Gardner¹, E.R. Atwill³, B.A. Byrne¹, S. Jang¹, M. Harris², J. Ames², D. Jessup², D. Paradies⁴, K. Worcester⁵, A. Melli¹ and P.A. Conrad¹.

(1) School of Veterinary Medicine, University of California, Davis, CA 95616, USA

(2) California Department of Fish and Game, Marine Wildlife Veterinary Care and Research Center, Santa Cruz, CA 95060, USA

(3) Veterinary Medicine Teaching and Research Center, University of California, Davis, Tulare, CA 93274, USA

(4) Bay Foundation of Morro Bay, Los Osos, CA 93412 USA

(5) Central Coast Regional Water Quality Control Board, San Luis Obispo, CA 93401 USA

Corresponding author:

Woutrina Miller, VM:PMI, 1126 Haring Hall, One Shields Ave, Davis, CA 95616, +1 530 219-1369 ph, +1 530 752-3349 f, wasmith@ucdavis.edu.

Submitted: 6 November 2005

Resubmitted: 23 February 2006

Running head: Coastal invertebrate bacterial ecology

Abstract

The coastal ecosystems of California are highly utilized by humans and animals, but the ecology of fecal bacteria at the land-sea interface is not well understood. This study evaluated the distribution of potentially pathogenic bacteria in invertebrates from linked marine, estuarine, and freshwater ecosystems in central California. A variety of filter-feeding clams, mussels, worms, and crab tissues were selectively cultured for *Salmonella* spp., *Campylobacter* spp., *Escherichia coli*-O157, *Clostridium perfringens*, *Plesiomonas shigelloides*, and *Vibrio* spp. A longitudinal study assessed environmental risk factors for detecting these bacterial species in sentinel mussel batches. Putative risk factors included mussel collection near higher risk areas for livestock or human sewage exposure, adjacent human population density, season, recent precipitation, water temperature, water type, bivalve type, and freshwater outflow exposure. Bacteria detected in invertebrates included *Salmonella* spp., *Clostridium perfringens*, *Plesiomonas shigelloides*, *Vibrio cholerae*, *Vibrio parahaemolyticus*, and *Vibrio alginolyticus*. Overall, 80% of mussel batches were culture-positive for at least one of the bacterial species, though the pathogens *Campylobacter*, *E. coli*-O157, and *Salmonella* were not detected. Many of the same bacterial species were also cultured from upstream estuarine and riverine invertebrates. Exposure to human sewage sources, recent precipitation, and water temperature were significant risk factors for bacterial detection in sentinel mussel batches. These findings are consistent with the hypothesis that filter-feeding invertebrates along the coast concentrate fecal bacteria flowing from land to sea, and show that the relationships between anthropogenic effects on coastal ecosystems and the environmental niches of fecal bacteria are complex and dynamic

43 *Introduction*

44 The ecology of fecal bacteria entering nearshore aquatic environments along the California
45 coast is not well understood, but is gaining in importance as human populations utilizing these
46 ecosystems grow (8, 37, 40, 43). Environmental loading of fecal byproducts from humans and
47 their associated animals is significant and can affect the quality of water and food resources in
48 coastal ecosystems (13, 24). Terrestrial sources of fecal waste entering aquatic ecosystems
49 include animal-derived fecal matter carried in storm runoff, and human sources such as sewage
50 outfalls, leaky septic tanks, and boat discharges. Many of the fecal parasites, bacteria, and
51 viruses that have been detected in freshwater, estuarine, and marine environments are potential
52 pathogens to humans and animals, and their environmental niches are quite diverse (6, 36).

53 Our previous studies have investigated the epidemiology of the fecal protozoan parasites
54 *Cryptosporidium*, *Giardia*, and *Toxoplasma* species in coastal California ecosystems (8, 28-30,
55 32). All of these parasite life cycles involve fragile zoite stages that develop in the digestive
56 tracts of animal hosts, including in some cases humans, and sturdy oocyst or cyst stages that are
57 shed in the feces and can remain viable in the environment for prolonged periods of time (12, 39).
58 Investigation into the ecology of these parasites in aquatic ecosystems revealed that detection of
59 *Cryptosporidium* and *Giardia* spp. in bivalve shellfish was associated with exposure to
60 freshwater sources and was greatest during the wet season in California (30, 32). *Toxoplasma*
61 *gondii* in a marine mammal, the Southern sea otter (*Enhydra lutris nereis*), has also been
62 correlated with exposure to areas of high freshwater outflow into the nearshore marine
63 environment (28). While the life cycles of fecal bacteria are quite different than fecal protozoa,
64 some of the same risk factors may apply to the transport of fecal bacteria from land to sea.

Fecal bacteria entering aquatic ecosystems in coastal California may have a more complex ecological niche because unlike the protozoans that are only known to multiply within animal cells, bacteria can multiply in the environment as well as within animal hosts (18, 44). This suggests that once fecal bacteria enter the aquatic environment, they may be able to establish long-term reservoirs, depending on each species' ability to acclimate and tolerate a range of environmental conditions. The bacteria in this study include primarily terrestrial fecal bacteria such as *Salmonella* and *Campylobacter* spp., as well as fecal bacteria well-adapted to aquatic environments such as *Vibrio* and *Plesiomonas* spp., all of which can act as pathogens under the appropriate host, microbe, and environmental conditions (2, 5, 18, 40, 44).

Filter-feeding invertebrates play an important role in the ecology of aquatic pathogens because they concentrate microbes from the surrounding waters (4, 16, 27). As a result, filter-feeders can act as bioindicators of the microbial diversity present in aquatic ecosystems, as well as food items that provide concentrated microbial doses to the animals and humans that consume them. The goal of this study was to investigate the distribution of potentially pathogenic bacteria among aquatic invertebrates in marine, estuarine, and freshwater ecosystems along the central California coast. Our primary hypothesis was that if fecal pollution is flowing from land to sea, then detection of fecal bacteria in invertebrates may be greatest near sites designated as higher risk for livestock runoff or human sewage exposure, compared to sites designated as lower risk for these sources. Based on our previous protozoal studies, we also hypothesized that high freshwater outflow exposure and wet season sampling would be associated with greater detection of fecal bacteria in bivalves.

Methods

88 *Study Design.* Bacteria selected for this study included *Salmonella* spp., *Campylobacter*
89 spp., *Escherichia coli*-O157, *Clostridium perfringens*, *Vibrio* spp., and *Plesiomonas shigelloides*.

90 Three approaches were taken to evaluating their distribution among invertebrates along the
91 central California coast. First, a longitudinal study was set up to deploy and collect sentinel
92 mussel batches for quarterly testing during the dry (June-November) and wet (December-May)
93 seasons along the central California coast. Figure 1 shows the various mussel sites sampled over
94 the course of the three year study, 2001-2004. Mussels were initially collected from a pristine
95 site near Bodega Bay, California, from which the target bacteria were not detected at any point
96 during the study, and were deployed at three sites considered at higher risk for human sewage
97 exposure, three sites at higher risk for livestock runoff, and three sites distant to these fecal
98 sources (32). Mussels were collected quarterly as batches of 30 and when transplanted mussels
99 were not available, resident mussels from the study sites were collected instead. Data on
100 environmental risk factors that could be associated with fecal pathogen pollution flowing from
101 terrestrial to aquatic ecosystems were also collected for each study site and sampling time.

102 Second, a comparative study was conducted of multiple invertebrate species collected from
103 muddy and sandy habitats near Moss Landing (Monterey Bay) and Morro Bay, California.

104 Clams, innkeeper worms, and mussels were sampled from a muddy estuarine habitat on the same
105 day that clams, sand crabs, and mussels were collected from a nearby sandy habitat just outside
106 the estuary during the wet season of 2002-2003. The target sample size was again 30 per species.

107 Third, freshwater clams (*Corbicula fluminea*) were deployed in the San Lorenzo and Salinas
108 Rivers, two rivers with high historic coliform counts along the central coast ([http://www.water](http://www.waterboards.ca.gov/centralcoast/BasinPlan/Documents/3BactiObjsStaffReport05-04-04.doc)
109 [boards.ca.gov/centralcoast/BasinPlan/ Documents/3BactiObjsStaffReport05-04-04.doc](http://www.waterboards.ca.gov/centralcoast/BasinPlan/Documents/3BactiObjsStaffReport05-04-04.doc)), both of
110 which feed into the Monterey Bay as shown in Figure 1. Clams were collected in batches of 30,

111 once during the dry season and again during the wet season for two consecutive years, 2002-
112 2004. Bivalve collection and deployment methods were in accordance with long-term coastal
113 monitoring protocols (30).

114

115 *Invertebrate Sampling.* Surf mussels (*Mytilus californianus*) and bay mussels (*Mytilus*
116 *edulis*) were used in the longitudinal mussel study. Invertebrate species sampled from the
117 estuaries near Moss Landing and Morro Bay, CA included fat innkeeper worms (*Urechis caupo*),
118 gaper clams (*Tresus nuttalli*), Washington clams (*Saxidomus nuttalli*), bent-nose clams (*Macoma*
119 spp.), and mussels (*Mytilus* spp.). Sampling from the sandy habitats outside the estuaries
120 included Pismo clams (*Tivela stultorum*), mussels (*Mytilus* spp.) and two types of sand crabs
121 (*Emerita analoga* and *Blepharipoda occidentalis*). Freshwater clams (*Corbicula fluminea*) were
122 deployed for invertebrate testing in the San Lorenzo and Salinas Rivers.

123 Invertebrates were collected as batches of 30, of which six individuals were pooled per
124 culture batch from each site and timepoint. Invertebrates were transported on ice to laboratories
125 at the University of California, Davis (UCD), or to the Marine Wildlife Veterinary Care and
126 Research Center in Santa Cruz, CA, for dissection within 24 hours of collection. Hemolymph,
127 gill, and digestive gland tissues were collected as described (30, 32), and cultured for bacteria
128 using selective media at UCD. The best invertebrate tissue to culture was determined by
129 comparing the results of culturing hemolymph, gill and digestive gland tissues from the same
130 invertebrates in the first year, with only digestive gland cultured in subsequent years.

131

132 *Bacterial Culture.* A 1-2 g aliquot of invertebrate tissues was macerated with a sterile swab
133 and saline in a microcentrifuge tube before plating onto selective media according to established

134 protocols (34). Cultures were enriched for *Salmonella* and *Vibrio* spp. for 24 hrs using selenite
135 broth and alkaline peptone water before plating onto Xylose Lactose Tergitol 4 (XLT4) and
136 Thiosulfate Citrate Bile Sucrose (TCBS) agar, respectively. *Campylobacter* cultures utilized
137 selective media containing Cefoperazone Vancomycin and Amphotericin B (Campy CVA) in a
138 microaerophilic environment. *Escherichia coli*-O157 cultures were performed on Cefixime
139 Potassium Tellurite Sorbitol MacConkey (CT SMAC) plates. *Clostridium perfringens* was
140 identified using Egg Yolk Agar (EYA) under anaerobic conditions. *Plesiomonas* was identified
141 from MacConkey plates. All incubations were performed at 37°C. Morphologic appearance and
142 biochemical testing was used to initially identify the isolates, with further characterization using
143 specific antiserum, API 20E strips, serotyping, and sequence analysis of the 16S rRNA gene.

145 *Risk Factors.* Environmental factors that could be associated with bacterial detection in
146 mussels were identified and the data assimilated for all study sites and timepoints. Categorical
147 data on fecal risk category was defined as lower risk for fecal pollution if sites were more than 5
148 km away from known fecal sources, as higher risk for livestock fecal sources if sites were less
149 than 5 km from livestock runoff, and as higher risk for human sewage exposure if sites were less
150 than 5 km from a sewage outfall or historic septic leakage. The distance of 5 km was selected
151 based on biologic plausibility in a previous study (32) and in consultation with local resource
152 managers who identified known fecal input sources within a 5 km radius of the sampling site.
153 Season was categorized by dry (June-November) or wet (December-May) season for each
154 collection. Recent precipitation was recorded as present or absent for the past day and past week.
155 Water type at each collection site was defined as estuarine or marine. Bivalve type was recorded
156 as resident or transplanted for each mussel batch. Freshwater outflow exposure data, defined as

low = <10 million L, medium = 10-100 million L, high = >100 million L of freshwater exposure in the past day, were available for each study site and mussel collection date from our previous *Cryptosporidium* study (32). In addition, human densities of the adjacent coastline were estimated using 2000 census data (<http://www.census.gov/main/www/cen2000.html>) and water temperatures for each mussel collection date were obtained from National Oceanic and Atmospheric Administration records (<http://co-ops.nos.noaa.gov/>).

Statistical Analysis. McNemar's χ^2 test for paired data was used to evaluate whether significant differences in culture success could be detected among hemolymph, gill, and digestive tissues from the same bivalves, as well as comparing culture results from mud and sand invertebrates collected at the same estuaries. For the mussel risk factor analysis, univariate and then multivariate logistic regression models were created to quantify the strength of association between environmental variables and detection of the enteric bacteria of interest. Logistic regression is a powerful tool for simultaneously assessing the relative contributions of multiple risk factors that may be associated with a dichotomous outcome of interest such as the presence or absence of a bacterial species. The odds ratio produced for each risk factor is interpreted as the probability that the outcome of interest will occur in samples with the risk factor, relative to the probability of the outcome occurring in samples without the risk factor (or referent category). To minimize sparse data problems, models were only created for bacteria that were detected more than five times over the three year study (19). No study sites were within 5 km of each other, and only one site (the most downstream site) per estuary was included. A cluster effect was used in all logistic regression models, to adjust for repeated sampling at the same sites over

179 time. All statistical analyses were performed using Stata software (Stata Corp, College Station,
180 TX, USA). Significant P values were defined as $P < 0.1$.

181

182 **Results**

183 *Tissue Comparison Study.* During the first year of bivalve collections, the hemolymph,
184 gill, and digestive gland tissues from the same animals ($n=110$) were cultured to determine the
185 optimal tissue for long-term study. There was no significant difference between the frequency of
186 isolation of enteric bacteria from hemolymph or gill tissues ($P=0.4$). However, digestive gland
187 tissues significantly outperformed both hemolymph and gill tissues ($P < 0.001$). Based on these
188 results, only digestive gland was cultured for the remainder of the study.

189

190 *Mussel Study.* The three year mussel study involved deploying and collecting mussels quarterly
191 from sites ranging south from Bodega Bay to Morro Bay, CA as shown in Figure 1. Forty-six
192 mussel batches met the study inclusion criteria for statistical analysis, with each study site
193 sampled at least three times during the study. Table 1 shows the bacterial prevalence among the
194 46 batches in the dry season, wet season, and in total. Overall, 80% of mussel batches during the
195 dry and wet seasons were positive for at least one of the bacterial species, though no
196 *Campylobacter*, *E.coli*-O157, or *Salmonella* spp. were detected. *Plesiomonas shigelloides* was
197 detected in 4% of batches collected during the dry season but not during the wet season. *Vibrio*
198 *parahaemolyticus*, *V. cholerae* non-O1, and *V. alginolyticus* were detected in up to 48% of
199 mussel batches collected during the dry season and up to 26% of mussel batches collected during
200 the wet season. In contrast, *C. perfringens* was detected more often in the wet season (68%) than
201 in the dry season (41%).

Univariate analyses evaluated the strength of association between individual environmental variables and a mussel batch culturing positive for each bacterial group. The odds ratios and associated P values for factors significantly associated with detection of *V. cholerae*, *V. alginolyticus*, *C. perfringens*, and 'Any bacterial species' outcomes are given in Table 2. The other individual bacterial species were not included due to the sparse nature of the data. *Vibrio cholerae* was not significantly associated with any of the environmental variables. A mussel batch was significantly less likely to test positive for *V. alginolyticus* during the wet season or if there was precipitation in the day or week preceding mussel collection. A batch was more likely to have *V. alginolyticus* isolated if the water temperature was greater than 12°C at the time of sampling or if the study site was marine as compared to estuarine. Detecting *C. perfringens* in mussel batches was significantly more likely to occur during the wet season, if there was not precipitation in the day preceding mussel collection, or if the water temperature was less than 12°C. Mussel batches in which any of the bacteria were cultured were significantly associated with the higher risk category for exposure to human sewage, and with not having precipitation in the day or week preceding mussel collections. Human density and bivalve type were not significantly associated with any of the bacterial outcomes.

Forward-stepping multivariate logistic regression was used to examine multiple environmental variables simultaneously against the outcome of a batch culturing positive for the bacteria of interest. Table 3 shows the three models for which significant explanatory variables were found. Culturing *V. alginolyticus* from mussel batches collected within 24 hours of a precipitation event was 5 times less likely (odds ratio = 0.2) than mussel batches in which no precipitation occurred in the preceding day ($P < 0.01$). *Clostridium perfringens* was also less likely to be detected in mussel batches collected within a day following a precipitation event

225 (P=0.07), and was associated with water temperatures less than 12°C (P=0.02). Detecting any of
 226 the targeted bacterial species in a mussel batch was significantly associated with no precipitation
 227 in the day preceding mussel collection (P<0.01), and was 39 times greater if the mussel batch
 228 was collected from a site considered higher risk for human sewage exposure compared to sites
 229 considered lower risk for human sewage or livestock waste exposure (P=0.01).

230

231 *Comparative Invertebrate Study.* The results of culturing pools of six invertebrates from
 232 muddy and sandy habitats near two fecal-impacted estuaries in the wet season of 2002-2003 are
 233 shown in Table 4. Bacterial pathogens were detected in the innkeeper worms, bivalves, and sand
 234 crabs that live in the sediment, as well as in mussels that were suspended higher up in the water
 235 column. *Salmonella* Typhimurium was detected in Pismo clams and *S. Heidelberg* in fat
 236 innkeeper worms at the Moss Landing site. *Vibrio parahaemolyticus* was cultured from Pismo
 237 clams and from at the Morro Bay site. *Vibrio alginolyticus* and *Clostridium perfringens* were
 238 detected in a variety of invertebrate species at both sites. Invertebrate pools (n=44) from the
 239 muddy sites inside the estuaries were more often positive for enteric bacteria than invertebrates
 240 collected at the sandy sites just outside the estuaries (P<0.01).

241

242 *Freshwater Clam Study.* The prevalence of each bacterial species cultured from pools of six
 243 freshwater clams collected during the 2002-2004 dry and wet seasons is shown in Table 5.
 244 *Vibrio cholerae* *V. alginolyticus*, and *C. perfringens* were only detected during the dry seasons.
 245 Downstream clam batches were more often culture-positive for bacteria than upstream batches.
 246 No *Campylobacter*, *E.coli*-O157, *Plesiomonas*, *Salmonella*, or other *Vibrio* spp. were detected.
 247 During the first wet season sampling, no clams were found alive at the lower San Lorenzo River

248 site and both Salinas River sites. Clams were repeatedly deployed at a third Salinas River site but
249 the clam batches were consistently missing at each sampling timepoint. Sites and time periods
250 for which no clams were cultured were excluded from further analysis.

251

252 ***Discussion***

253 The goal of this study was to undertake the first multiyear investigation of the distribution of
254 potentially pathogenic bacteria among aquatic invertebrates in linked marine, estuarine, and
255 freshwater ecosystems along the Pacific coast of North America. *Campylobacter* and *E.coli*-
256 O157 were not detected in the study, *Salmonella* and *Plesiomonas shigelloides* were only rarely
257 detected, and *Clostridium perfringens* and *Vibrio* spp. were commonly detected. We had
258 hypothesized that if biologic pollution due to fecal bacteria is flowing from land to sea, then
259 detection of these organisms in invertebrates would be greatest near sites at higher risk for
260 exposure to livestock or human feces, compared to sites designated as lower risk for these
261 sources. This hypothesis was supported by our risk factor analysis, in which detection of any of
262 these bacteria was greatest in invertebrates collected near areas at higher risk for exposure to
263 human sewage. However, no significant association was observed between detecting the
264 targeted bacteria and invertebrates collected from areas at higher risk for exposure to livestock
265 runoff, nor were either of the higher risk fecal categories significant risk factors in the logistic
266 regression models for individual bacteria. Study sites considered at higher risk for human
267 sewage exposure included sites near human sewage treatment outfalls, as well as sites influenced
268 by leaking septic tanks. In addition, other non-point sources such as urban runoff could be
269 transporting human and animal fecal bacteria from land to sea in these higher risk areas but could
270 not be assessed in this study. Detecting potentially pathogenic bacteria in invertebrates exposed

271 to human sewage could be due to increased pathogen input into the marine ecosystem with
272 human waste, but could also be due to increased nutrient inflow that could make environmental
273 conditions more favorable for bacteria to grow. Other studies have previously detected
274 associations between compromised water quality in nearshore marine ecosystems and human
275 sewage inputs (24, 25, 35).

276 Based on our recent studies, we also hypothesized that high freshwater outflow exposure and
277 wet season sampling would be associated with detection of fecal bacteria in bivalves (11, 28, 32).
278 In this study, high freshwater outflow exposure was not significantly associated with detecting
279 bacteria in mussels, and while *C. perfringens* was detected most often in the wet season, *Vibrio*
280 spp. and *P. shigelloides* were detected most often in the dry season. These findings suggest that
281 the ecology of bacteria in coastal ecosystems is quite different than the ecology of fecal-borne
282 protozoa. All of the bacteria in this study can be shed in the feces of humans and animals, but
283 unlike the protozoa, many bacteria also survive and grow in environmental reservoirs when the
284 conditions are favorable (9, 18). In other studies, *Vibrio* and *Plesiomonas* spp. have been
285 detected most often during the warmer temperatures of summer and under certain salinity ranges
286 (6, 18, 44). Furthermore, *Vibrio* spp. have been shown to play a role in chitin degradation,
287 biofilm formation, and microbial competence in aquatic ecosystems, again highlighting their
288 specialized ecologic niche in aquatic ecosystems (2).

289 *Clostridium perfringens* was not associated with the same environmental parameters as the
290 other bacteria targeted in our study or others (4, 9). Several studies have detected *C. perfringens*
291 throughout the year under a variety of conditions, and it has been utilized as an indicator for fecal
292 source tracking studies (1, 25, 35). *Clostridium perfringens* is an anaerobic, spore-forming
293 bacteria shed in the feces of a variety of animals and humans, with some genotypes associated

294 with clinical disease (34). The finding that both *Vibrio* spp. and *C. perfringens* were
295 significantly associated with mussels collected when no precipitation had occurred in the
296 previous day suggests that rain may influence the bacteria-invertebrate interactions, possibly by
297 altering the bacterial concentration in surface waters, and/or the invertebrate feeding and
298 depuration dynamics (33).

299 Several types of *Vibrio* spp. were detected in this study. *Vibrio cholerae* is the most
300 infamous of the *Vibrios*, and while none of the epidemic-causing O1 strains were detected in this
301 study, even non-O1 *V. cholerae*s have been associated with clinical disease in humans in
302 California and elsewhere (18, 22, 36). *Vibrio parahaemolyticus* is another potentially pathogenic
303 strain that can cause enteritis, and *V. alginolyticus* is the most widespread environmental strain,
304 causing gastroenteritis or wound infections under opportune conditions (3, 18). The current
305 study did not test bacterial isolates for virulence-associated factors to more definitively evaluate
306 their potential pathogenicity, nor did it quantitate bacterial concentrations, another important
307 consideration when specifically evaluating microbial disease potential (5).

308 *Vibrio cholerae* was first reported along the California coast in the 1980's (21-23). The
309 Kenyon et al. (22, 23) studies were undertaken as a follow-up to a human case of *V. cholerae*
310 non-O1 gastroenteritis, and the highest *V. cholerae* levels were detected in water samples during
311 the summer, at times when coliform counts exceeded the legal limit of 1,000 per 100 ml seawater.
312 The Kaysner et al. (21) study tested water, sediment, and shellfish samples for *V. cholerae* from
313 24 estuaries along the U.S. Pacific coast. *Vibrio cholerae* non-O1 was detected in Washington,
314 Oregon, and California, with *V. cholerae* O1 Inaba reported in three water samples from Morro
315 Bay, CA. Predominantly non-O1 *V. cholerae* strains have also been reported on the gulf and east
316 coasts of the United States (20, 46).

Two serotypes of *Salmonella* were detected in invertebrates during the wet season collections. *Salmonella* Heidelberg was cultured from fat innkeeper worms living in the muddy estuary floor and *S. Typhimurium* was detected in Pismo clams from the sandy beach just outside the estuary mouth. Both of these *Salmonella* serotypes have been associated with clinical disease and are shed in the feces of a variety of hosts (17, 38, 45). Because *Salmonella* spp. were not cultured from mussels collected at the same time as the benthic invertebrates, it is uncertain whether the *Salmonella* were passing through in the water or were already present in the benthic sediment when filtered by the invertebrates. For most of the other bacterial species, suspended mussels and adjacent benthic invertebrates were culture-positive at the same sampling point, suggesting that the bacteria were passing through in the water column.

The comparative invertebrate and freshwater clam surveys showed that many of the same bacterial species isolated from marine mussels were also detectable in upstream invertebrates, though increased sample sizes and source tracking are needed to evaluate these associations statistically. Both estuaries in the study are at the interface between human activities and natural ecosystems, draining watersheds impacted by a wide range of agricultural and urban activities. Both have had problems with pollution and water quality, though invertebrates from both areas are still consumed by humans and marine mammals, such as the federally-listed threatened Southern sea otter (<http://www.waterboards.ca.gov/centralcoast/BasinPlan/Documents/3BactiObjsStaffReport05-04-04.doc>). More invertebrates were positive for bacteria from the muddy sites inside the estuaries than from the sandy sites just outside the estuaries. This finding is consistent with sources of bacteria coming from inland waterways and draining out into the ocean. However, it is also possible that the different environmental conditions inside the estuary are simply more favorable for bacterial survival than outside the estuary, allowing for bacteria

that enter the estuary on the incoming tide to find environmental niches in muddy habitat that aren't available in sandy habitat.

Mussels (*Mytilus* spp.) and freshwater clams (*Corbicula* spp.) were used for outplanting as sentinel bivalves for specific reasons. First, they already existed in the ecosystems of interest along the California coast, and they had been used for years by coastal monitoring programs to study pesticide and metal pollutants in coastal ecosystems (41). Second, the inherent filter-feeding activities of bivalves make them a natural concentrating mechanism that is easier to process in the laboratory than large volumes of water. Other invertebrates worth considering for sentinel studies include fat innkeeper worms in muddy habitats, that were recently shown to retain much higher levels of the algal neurotoxin domoic acid than other invertebrates along the California coast (<http://seafloor.csUMB.edu/publications/capstones/goldbergthesis.pdf>), and sand crabs that live on sandy beaches where mussels are not always found, and have been shown to retain high levels of domoic acid in recent studies (14).

The culture methods used in this study were chosen because they are widely accepted and cost-efficient compared to molecular methods. The use of selective media allowed for efficient sample screening of a large number of samples, followed by further biochemical and molecular confirmation techniques on a more limited sample set. As has been shown in prior studies (10, 30, 31), digestive gland was the most sensitive tissue for screening purposes. It is possible that some samples falsely tested negative by culture in our study due to differences between the selective culture conditions and natural environmental conditions, or if the bacteria were in a viable but non-culturable state (15).

In conclusion, this study assessed a number of invertebrate species for potentially pathogenic bacteria present in coastal California ecosystems. The increased detection of bacteria in

363 invertebrates exposed to human sewage sources suggests that anthropogenic changes to the
364 nearshore marine ecosystem may have significant effects on the ecology of fecal bacteria at the
365 land-sea interface. Other studies support this association (11, 13, 26), making research on
366 methods to minimize the impacts of humans and their associated animals on nearshore
367 ecosystems important for long-term sustainability and health (1, 7, 42).

368

369 *Acknowledgements*

370 This study was supported in part by the National Sea Grant College Program under NOAA
371 grant NAO6RG0142 Project R/CZ-180, through the California Sea Grant College Program; and
372 in part by the California State Resources Agency and the California Department of Fish & Game.
373 This study was also supported by the University of California Water Resources Program,
374 Wildlife Health Center, Center for Food Animal Health, and the NIH NIGMS Professors of the
375 Future Program. The assistance of Gary Ichikawa at the California State Mussel Watch Program,
376 Bryn Phillips at the Granite Canyon Marine Lab, Paul Olin at the California Sea Grant College
377 Program, David Lewis at the University of California Cooperative Extension, the California Fish
378 and Game staff, local stakeholders, and University of California students is appreciated.

379

380 *References*

- 381 1. Barrett, EC, Sobsey, MD, House, CH, White, KD (2001) Microbial indicator removal in onsite
382 constructed wetlands for wastewater treatment in the southeastern U.S. Water Sci Technol
383 44:177-182.
- 384 2. Bartlett, DH, Azam, F (2005) Chitin, cholera, and competence. Science 310:1775-1777.

- 385 3. Blake, PA (1983) Vibrios on the half shell: what the walrus and carpenter didn't know. Ann
386 Intern Med 99:558-559.
- 387 4. Burkhardt, W, Watkins, WD, Rippey, SR (1992) Seasonal effects on accumulation of
388 microbial indicator organisms by *Mercenaria mercenaria*. Appl Environ Microbiol 58:826-
389 831.
- 390 5. Butt, AA, Aldridge, KE, Sanders, CV (2004) Infections related to the ingestion of seafood,
391 Part I: viral and bacterial infections. Lancet 4:201-212.
- 392 6. Chowdhury, MA., Miyoshi, S, Yamanaka, H, Shinoda, S (1992) Ecology and distribution of
393 toxigenic *Vibrio cholerae* in aquatic environments in a temperate region. Microbios 72:203-
394 213.
- 395 7. Collins, R, Rutherford, K (2004) Modelling bacterial water quality in streams draining
396 pastoral lands. Water Res 38:700-712.
- 397 8. Conrad, PA, Miller, MA, Kreuder, C, James, ER, Mazet, J, Dabritz, H, Jessup, DA, Gulland,
398 F, Grigg, ME (2005) Transmission of *Toxoplasma*: clues from the study of sea otters as
399 sentinels of *Toxoplasma gondii* flow into the marine environment. Int J Parasitol Sep 10 [Epub
400 ahead of print].
- 401 9. Desmarais, TR, Solo-Gabrielle, HM, Palmer, CJ (2002) Influence of soil on fecal indicator
402 organisms in a tidally influenced subtropical environment. Appl Environ Microbiol 68:1165-
403 1172.

- 1
2
3 404 10. Dore, WJ, Lees, DN (1995) Behavior of *Escherichia coli* and male-specific bacteriophage in
4
5 405 environmentally contaminated bivalve molluscs before and after depuration. Appl Environ
6
7 406 Microbiol 61:2830-2834.
8
9
10
11 407 11. Dwight, RH, Semenza, JC, Baker, DB, Olson, BH (2002) Association of urban runoff with
12
13 408 coastal water quality in Orange County, California. Water Environ Res 74:82-90.
14
15
16
17 409 12. Fayer, R, Ed (1997) *Cryptosporidium* and Cryptosporidiosis, CRC Press, Boston.
18
19
20
21 410 13. Fayer, R, Dubey, JP, Lindsay, DS (2004) Zoonotic protozoa: from land to sea. Trends
22
23 411 Parasitol 20:531-536.
24
25
26
27 412 14. Ferdin, ME, Kvitek, RG, Bretz, CK, Powell, CL, Doucette, GJ, Lefevre, KA, Coale, S,
28
29 413 Silver, MW (2002) *Emerita analoga* (Stimpson)-possible new indicator species for the
30
31 414 phycotoxin domoic acid in California coastal waters. Toxicon 40:1259-1265.
32
33
34
35 415 15. Garcia-Armison, T, Servais, P (2004) Enumeration of viable *E. coli* in rivers and wastewaters
36
37 416 by fluorescent in situ hybridization. J Microbiol Methods 58:269-279.
38
39
40
41 417 16. Graczyk, TK, Conn, DB, Marcogliese, DJ, Graczyk, H, De Lafontaine, Y (2002)
42
43 418 Accumulation of human waterborne parasites by zebra mussels (*Dreissena polymorpha*) and
44
45 419 Asian freshwater clams (*Corbicula fluminea*). Parasitol Res 89:107-112.
46
47
48
49 420 17. Guerin, MT, Marin, SW, Darlington, GA, Rajic, A (2005) A temporal study of *Salmonella*
50
51 421 serovars in animals in Alberta between 1990 and 2001. Can J Vet Res 69:88-99.
52
53
54
55 422 18. Holmberg, SD (1988) *Vibrios* and *Aeromonas*. Infect Dis Clin North Am 2:655-676.
56
57
58
59
60

- 1
2
3 423 19. Hosmer, DW, Lemeshow, S (2000) Applied Logistic Regression, 2nd Ed, Ch. 8, John Wiley
4
5
6 424 & Sons, New York, New York.
7
8
9 425 20. Kaper, J, Lockman, H, Colwell, RR, Joseph, SW (1979) Ecology, serology, and enterotoxin
10
11 426 production of *Vibrio cholerae* in Chesapeake Bay. Appl Environ Microbiol 37:91-103.
12
13
14
15 427 21. Kaysner, CA, Abeyta, C, Wekell, MM, DePaolo, A, Stott, RF, Leitch, JM (1987) Incidence
16
17 428 of *Vibrio cholerae* from estuaries of the United States west coast. Appl Environ Microbiol
18
19 429 53:1344-1348.
20
21
22
23 430 22. Kenyon, JE, Gillies, DC, Piexoto, DR, Austin, B (1983) *Vibrio cholerae* (non-O1) isolated
24
25 431 from California coastal waters. Appl Environ Microbiol 46:1232-1233.
26
27
28
29 432 23. Kenyon, JE, Piexoto, DR, Austin, B, Gillies, DC (1984) Seasonal variation in numbers of
30
31 433 *Vibrio cholerae* (non-O1) isolated from California coastal waters. Appl Environ Microbiol
32
33 434 47:1243-1245.
34
35
36
37 435 24. Kim, JH, Grant, SB, McGee, CD, Sanders, BF, Largier, JL (2004) Locating sources of surf
38
39 436 zone pollution: a mass budget analysis of fecal indicator bacteria at Huntington Beach,
40
41 437 California. Environ Sci Technol 38:2626-2636.
42
43
44
45
46 438 25. Lipp, EK, Farrah, SA, Rose, JB (2001) Assessment and impact of microbial fecal pollution
47
48 439 and human enteric pathogens in a coastal community. Mar Pollut Bull 42:286-293.
49
50
51
52 440 26. Mallin, MA, Williams, KE, Esham, EC, Lowe, RP (2000) Effect of human development on
53
54 441 bacteriological water quality in coastal watersheds. Ecol Appl 10:1047-1056.
55
56
57
58
59
60

- 442 27. Marino, A, Lombardo, L, Fiorentino, C., Orlandella, B, Monticelli, L, Nostro, A, Alonzo, V
443 (2005) Uptake of *Escherichia coli*, *Vibrio cholerae* non-O1 and *Enterococcus durans* by, and
444 depuration of mussels (*Mytilus galloprovincialis*). Int J Food Microbiol 99:281-286.
- 445 28. Miller, MA, Gardner, IA, Kreuder, C, Paradies, DM, Worcester, KR, Jessup, DA, Dodd, E,
446 Harris, MD, Ames, JA, Packham, AE, Conrad, PA (2002) Coastal freshwater runoff is a risk
447 factor for *Toxoplasma gondii* infection of southern sea otters (*Enhydra lutris nereis*). Int J
448 Parasitol 32:997-1006.
- 449 29. Miller, MA, Grigg, ME, Kreuder, C, James, ER, Melli, AC, Crosbie, PR, Jessup, DA,
450 Boothroyd, JC, Brownstein, D, Conrad, PA (2004) An unusual genotype of *Toxoplasma gondii*
451 is common in California sea otters (*Enhydra lutris nereis*) and is a cause of mortality. Int J
452 Parasitol 34:275-284.
- 453 30. Miller, WA, Atwill, ER, Gardner, IA, Miller, MA, Fritz, HM, Hedrick, RP, Melli, AC,
454 Barnes, NM, Conrad, PA (2005) Clams (*Corbicula fluminea*) as bioindicators of fecal
455 contamination with *Cryptosporidium* and *Giardia* spp. in freshwater ecosystems in California.
456 Int J Parasitol 35:673-684.
- 457 31. Miller, WA, Gardner, IA, Atwill, ER, Leutenegger, CM, Miller, MA, Hedrick, RP, Melli,
458 AC, Barnes, NM, Conrad, PA (2005) Evaluation of methods for improved detection of
459 *Cryptosporidium* spp. in mussels (*Mytilus californianus*). J Microbiol Methods, In Press.
- 460 32. Miller, WA, Miller, MA, Gardner, IA, Atwill, ER, Harris, M, Ames, J, Jessup, D, Melli, A,
461 Paradies, D, Worcester, K, Olin, P, Barnes, N, Conrad, PA (2005) New genotypes and factors

- 1
2
3 462 associated with detection of *Cryptosporidium* detection in mussels (*Mytilus* spp.) along the
4
5
6 463 California coast. Int J Parasitol 35:1103-1113.
7
8
9 464 33. Mourino-Perez, RR, Worden, AZ, Azam, F (2003) Growth of *Vibrio cholerae* O1 in red tide
10
11 465 waters off California. Appl Environ Microbiol 69:6923-6931.
12
13
14
15 466 34. Murray, P.R., Ed (1995) *Manual of Clinical Microbiology*. ASM Press, Washington D.C.
16
17
18 467 35. Paul, JH, Rose, JB, Jiang, S, Kellogg, C, Shinn, EA (1995) Occurrence of fecal indicator
19
20 468 bacteria in surface waters and the subsurface aquifer in Key Largo, Florida. Appl Environ
21
22 469 Microbiol 61:2235-2241.
23
24
25
26
27 470 36. Rippey, S (1994) Infectious diseases associated with molluscan shellfish consumption. Clin
28
29 471 Microbiol Rev 7:419-425.
30
31
32
33 472 37. Schlundt, J, Toyofuku, H, Jansen, J, Herbst, SA (2004) Emerging food-borne zoonoses. Rev
34
35 473 Sci Tech 23:513-533.
36
37
38
39 474 38. Seepersadsingh, N, Adesiyun, AA, Seebaransigh, R (2004) Prevalence and antimicrobial
40
41 475 resistance of *Salmonella* spp. in non-diarrhoeic dogs in Trinidad. J Vet Med B Infect Dis Vet
42
43 476 Public Health 51:337-342.
44
45
46
47 477 39. Sinski, E (2003) Environmental contamination with protozoan parasite infective stages:
48
49 478 biology and risk assessment. Acta Microbio Pol 52S:97-107.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 479 40. Skelly, C, Weinstein, P (2003) Pathogen survival trajectories: an eco-environmental
4
5 480 approach to the modeling of human campylobacteriosis ecology. Environ Health Perspect
6
7 481 111:19-28.
8
9
10
11 482 41. Stephenson, MD, Martin, M, Tjeerdema, RS (1995) Long-term trends in DDT,
12
13 483 polychlorinated biphenyls, and chlordane in California mussels. Arch Environ Contam
14
15 484 Toxicol 28:443-450.
16
17
18
19 485 42. Tate, KW, Pereira, MDGC, Atwill, ER (2004) Efficacy of vegetated buffer strips for
20
21 486 retaining *Cryptosporidium parvum*. J Environ Qual 33:2243-2251.
22
23
24
25 487 43. Turbow, DJ, Osgood, ND, Jiang, SC (2003) Evaluation of recreational health risk in coastal
26
27 488 waters based on enterococcus densities and bathing patterns. Environ Health Perspect 111:598-
28
29 489 603.
30
31
32
33
34 490 44. Venkataswaren, K, Takai, T, Navarro, IM, Nakano, H, Hashimoto, H, Siebeling, RJ (1989)
35
36 491 Ecology of *Vibrio cholerae* non-O1 and *Salmonella* spp. and role of zooplankton in their
37
38 492 seasonal distribution in Fukuyama coastal waters, Japan. Appl Environ Microbiol 55:1591-
39
40 493 1598.
41
42
43
44
45 494 45. Vugia, DJ, Samuel, M, Farley, MM, Marcus, R, Shiferaw, B, Shallow, S, Smith, K, Angulo,
46
47 495 FJ, Emerging Infections Program FoodNet working Group (2004) Invasive *Salmonella*
48
49 496 infections in the United States, FoodNet, 1996-1999: incidence, serotype distribution, and
50
51 497 outcome. Clin Infect Dis 38:S149-S156.
52
53
54
55
56
57
58
59
60

- 1
2
3 498 46. Weber, JT, Levine, WC, Hopkins, DP, Tauxe, RV (1994) Cholera in the United States, 1965-
4
5
6 499 1991, risks at home and abroad. Arch Intern Med 154:551-556.
7
8
9 500

500 Table and Figure captions:

501

502 Table 1. Prevalence of fecal bacteria in mussel batches, 2001-2004.

503

504 Table 2. Univariate analysis of factors significantly associated with detection of enteric bacteria
505 in mussel batches.

506

507 Table 3. Multivariate analysis of factors significantly associated with detection of enteric
508 bacteria in mussel batches.

509

510 Table 4. Bacteria cultured from invertebrate species living in two muddy and sandy estuarine
511 habitats in the wet season, 2002-2003.

512

513 Table 5. Bacteria cultured from freshwater clams in two coastal California Rivers, 2002-2004.^a

514

515 Figure 1. Map of the central California coast showing invertebrate field sampling sites.

516

517

518

519

520

521

522

523

524 Table 1

<i>Bacterial species</i>	<i>Seasonal prevalence(%)</i>		<i>Overall prevalence (%) (n=46)</i>
	<i>Dry (n=27 batches)</i>	<i>Wet (n=19 batches)</i>	
<i>Campylobacter</i>	0	0	0
<i>E.coli-O157</i>	0	0	0
<i>Salmonella</i>	0	0	0
<i>Plesiomonas shigelloides</i>	4	0	2
<i>Vibrio parahaemolyticus</i>	11	5	9
<i>Vibrio cholerae</i>	26	11	20
<i>Vibrio alginolyticus</i>	48	26	39
<i>Clostridium perfringens</i>	41	68	52
Any of the above bacteria	81	79	80

525

526

527

Table 2

Univariate risk factors	<i>Vibrio cholerae</i>		<i>Vibrio alginolyticus</i>		<i>Clostridium perfringens</i>		<i>Any bacterial species</i>	
	Odds ratio	P value	Odds ratio	P value	Odds ratio	P value	Odds ratio	P value
Fecal risk category - lower	1	-	1	-	1	-	1	-
higher - livestock sources	3	0.2	0.5	0.4	1.4	0.7	2.6	0.4
higher - human sources	3.7	0.3	2.4	0.3	2.1	0.4	13.6	0.08 ^a
Season: Dry (June-November)	1	-	1	-	1	-	1	-
Wet (December-May)	0.3	0.2	0.4	<0.01 ^a	3.2	0.04 ^a	0.9	0.7
Precipitation in the past day: no	1	-	1	-	1	-	1	-
yes	0.4	0.3	0.3	0.1 ^a	0.3	0.1 ^a	0.1	0.01 ^a
Precipitation in the past week: no	1	-	1	-	1	-	1	-
yes	0.6	0.3	0.2	<0.01 ^a	0.9	0.8	0.3	0.02 ^a
Water temperature over 12°C: no	1	-	1	-	1	-	1	-
yes	1.7	0.5	3.2	0.07 ^a	0.3	0.1 ^a	0.8	0.6
Water type: estuarine	1	-	1	-	1	-	1	-
marine	0.5	0.5	3.8	<0.01 ^a	0.9	0.8	0.3	0.3
Freshwater outflow exposure: low	1	-	1	-	1	-	1	-
medium	1.8	0.4	1.6	0.5	1	1	0.5	0.4
high	1.8	0.4	0.5	0.5	1.7	0.5	0.5	0.5

^aP value <0.1 considered significant.

Table 3

<i>Bacterial species</i>	<i>Significant risk factor</i>	<i>Odds ratio</i>	<i>P value^a</i>
<i>Vibrio alginolyticus</i>	Precipitation in the past week: no	1	-
	yes	0.2	<0.01
<i>Clostridium perfringens</i>	Precipitation in the past day: no	1	-
	yes	0.15	0.07
	Water temperature over 12°C: no	1	-
	yes	0.17	0.02
Any bacterial spp.	Fecal risk category: lower risk	1	-
	higher risk - livestock sources	6.9	0.2
	higher risk - human sources	39.4	0.01
	Precipitation in the past day: no	1	-
	yes	0.04	<0.01

^aP value <0.1 considered significant.

541 Table 4

542

543

544

545

546

<i>Site</i>	<i>Host species</i>	<i>Salmonella</i> ^b	<i>Vibrio</i> <i>parahaemolyticus</i> ^b	<i>Vibrio</i> <i>alginolyticus</i> ^b	<i>Clostridium</i> <i>perfringens</i> ^b
Moss Landing (mud)	Innkeeper worms	+	-	+	+
	Gaper clams	-	-	+	+
	Washington clams	-	-	+	+
	Bent-nose clams	-	-	-	+
	Mussels	-	-	+	+
Moss Landing (sand)	Pismo clams	+	-	-	-
	Blepharipoda crabs	-	-	-	-
	Emerita crabs	-	-	+	-
Morro Bay (mud)	Innkeeper worms	-	-	-	-
	Bent-nose clams	-	-	+	+
	Mussels	-	+	+	+
Morro Bay (sand)	Pismo clams	-	+	+	+
	Blepharipoda crabs	-	-	+	-
	Emerita crabs	-	-	-	-
	Mussels	-	-	+	+

^aNo *Campylobacter* spp., *E.coli* O157:H7, *Plesiomonas shigelloides*, or other *Vibrio* spp. were detected.

^b + = culture positive batch; - = culture negative batch.

547

548

549

549 Table 5

550

551

<i>Riverine Study Site</i>	<i>Vibrio cholerae</i>			<i>Vibrio alginolyticus</i>			<i>Clostridium perfringens</i>		
	<i>D1</i> ^b	<i>D2</i>	<i>W2</i>	<i>D1</i>	<i>D2</i>	<i>W2</i>	<i>D1</i>	<i>D2</i>	<i>W2</i>
San Lorenzo Upper	-	-	-	-	-	-	-	-	-
San Lorenzo Middle	-	-	-	+	-	-	+	-	-
San Lorenzo Lower	-	+	-	+	-	-	-	-	-
Salinas Upper	-	+	-	-	-	-	-	-	-
Salinas Lower	-	+	-	+	-	-	-	-	-

^a No *Campylobacter*, *E.coli* -O157, *Plesiomonas*, *Salmonella*, other *Vibrio* spp. were detected.

^b D = Dry season, W = Wet season; 1 = Year 1, 2 = Year 2.

552

